

**XX GEOMETRICAL OLYMPIAD IN HONOUR OF  
I.F.SHARYGIN  
The correspondence round. Solutions**

1. (8, D.Shvetsov) Bisectors  $AI$  and  $CI$  meet the circumcircle of triangle  $ABC$  at points  $A_1, C_1$  respectively. The circumcircle of triangle  $AIC_1$  meets  $AB$  at point  $C_0$ ; point  $A_0$  is defined similarly. Prove that  $A_0, A_1, C_0, C_1$  are collinear.

**Solution.** Let  $A_1C_1$  meet  $AB$  at point  $A'$ . Then

$$\angle C_1A'A = (\sphericalangle AC_1 + \sphericalangle BA_1)/2 = (\sphericalangle AC_1 + \sphericalangle A_1C)/2 = \angle C_1IA.$$

Therefore  $A, I, A', C_1$  are concyclic and  $A'$  coincides with  $A_0$  (fig. 1). Similarly we obtain that  $A_1C_1$  passes through  $C_0$ .




Fig. 1.

2. (8, B.Frenkin) Three different collinear points are given. What is the number of isosceles triangles such that these points are their circumcenter, incenter and excenter (in some order)?

**Answer.** Two, if the medial point divide the segment between two remaining ones into two parts such that their ratio is not greater than 3, and three otherwise.

**Solution.** Let  $ABC$  be an isosceles triangle with vertex  $C$ ;  $O, I, I_c$  be its circumcenter, incenter and excenter (center of excircle touching  $AB$ ). Then  $A, B, I, I_c$  lie on the circle with diameter  $II_c$  centered at the midpoint  $W$  of arc  $AB$ . Hence if  $O, I, I_c$  are given, then we construct the triangle  $ABC$  finding the midpoint  $W$  of segment  $II_c$ , drawing the circles centered at  $O, W$  with radii  $OW, WI$  respectively, finding the common points  $A$  and  $B$  of these circles, and the reflection  $C$  of  $W$  about  $O$ . But  $I$  has to lie inside the circumcircle. This is always correct if  $O$  is one of two extreme points (then  $I$  is the medial point). And if  $O$  is the medial point, then the inequality  $OW > OI$  has to be correct, i.e.  $OI_c : OI > 3$ .

3. (8, K.Belskij) Let  $ABC$  be an acute-angled triangle, and  $M$  be the midpoint of the minor arc  $BC$  of its circumcircle. A circle  $\omega$  touches the side  $AB$ ,  $AC$  at points  $P$ ,  $Q$  respectively and passes through  $M$ . Prove that  $BP + CQ = PQ$ .

**Solution.** By the assumption  $\angle PAQ = (\sphericalangle PMQ - \sphericalangle PQ)/2 = \pi - \sphericalangle PQ = \pi - 2\angle PMQ$ , i.e.  $\angle PMQ = \angle APQ$ . Thus  $PM$  and  $QM$  bisect the angles  $BPQ$ ,  $CQP$  respectively, and  $M$  is the incenter of triangle  $APQ$ . Then constructing perpendiculars  $MX$ ,  $MY$  to  $AB$ ,  $AC$  respectively we obtain that  $PX = QY = PQ/2$ . Also since  $MB = MC$ , we have  $BX = CY$  (fig. 3). Therefore  $BP + CQ = PX + XB + QY - YC = PQ$ .




Fig. 3.

4. (8, L.Emelyanov) The incircle  $\omega$  of triangle  $ABC$  touches  $BC$ ,  $CA$ ,  $AB$  at points  $A_1$ ,  $B_1$  and  $C_1$  respectively,  $P$  is an arbitrary point on  $\omega$ . The line  $AP$  meets the circumcircle of triangle  $AB_1C_1$  for the second time at point  $A_2$ . Points  $B_2$  and  $C_2$  are defined similarly. Prove that the circumcircle of triangle  $A_2B_2C_2$  touches  $\omega$ .

**Solution.** Points  $A$ ,  $B_1$ ,  $C_1$  lie on the circle with diameter  $AI$ , where  $I$  is the incenter of  $ABC$ . Hence  $\angle IA_2A = \angle IA_2P = 90^\circ$ , and  $A_2$  lies on the circle with diameter  $IP$ , touching  $\omega$  at  $P$  (fig. 4). Points  $B_2$ ,  $C_2$  also lie on this circle.




Fig. 4.

5. (8, P.Pogosjan) Points  $A'$ ,  $B'$ ,  $C'$  are the reflections of vertices  $A$ ,  $B$ ,  $C$  about the opposite sidelines of triangle  $ABC$ . Prove that the circles  $AB'C'$ ,  $A'BC'$ , and  $A'B'C$  have a common point.

**Solution.** Let  $X$  be the second common point of circles  $AB'C'$  and  $A'BC'$ . Then  $\angle(XB', XC') = \angle(AB', AC') = 3\angle(AC, AB)$ . Similarly  $\angle(XC', XA') = 3\angle(BA, BC)$ . Therefore  $\angle(XB', XA') = 3\angle(AC, BC) = \angle(CB', CA')$ .

6. (8–9, A.Shekera) A circle  $\omega$  and two points  $A$ ,  $B$  of this circle are given. Let  $C$  be an arbitrary point on one of arcs  $AB$  of  $\omega$ ;  $CL$  be the bisector of triangle  $ABC$ ; the circle  $BCL$  meet  $AC$  at point  $E$ ; and  $CL$  meet  $BE$  at point  $F$ . Find the locus of circumcenters of triangles  $AFC$ .

**Answer.** A segment with the endpoint at the midpoint of arc  $ACB$ , forming the angle with  $AB$  equal to  $\pi/2 - \angle ACB$

**Solution.** Let  $O$  be the center of circle  $ACF$ . Then  $\angle AOF = 2\angle ACF = \angle ACB$  do not depend on  $C$ . Hence all triangles  $AOF$  are similar, and  $O$  is the image of  $F$  about the spiral similarity with center  $A$ . Also  $\angle ABF = \angle LBE = \angle LCE$  do not depend on  $C$ , thus  $F$  moves along a line. Therefore all points  $O$  also lie on a line. The angle between this line and  $BE$  equals  $\angle OAF = (\pi - \angle ACB)/2 = \pi/2 - \angle ABE$ , hence it is perpendicular to the reflection of  $AB$  about  $BE$  (fig. 6). When  $C$  tends to  $B$ ,  $O$  tends to the midpoint of arc  $ACB$ , and when  $C$  tends to  $A$ ,  $F$  tends to the tangent to the circumcircle  $ABC$  at  $A$ . Therefore the required locus is the segment given in the answer.




Fig. 6.

7. (8–9, B.Frenkin) Restore a bicentral quadrilateral if two opposite vertices and the incenter are given.

**Solution.** Let the incenter  $I$  of bicentral quadrilateral  $ABCD$  lie inside triangle  $ABC$ . Then  $\angle AIC = \angle ABC + \angle IAB + \angle ICB = \angle ABC + \pi/2$ . Thus if  $A, C, I$  are given, we can find the value of angle  $B$  and construct the circumcircle centered at  $O$ . Now the line  $OI$  passes through the common point  $L$  of diagonals of  $ABCD$ , the midpoints  $M, N$  of  $AC, BD$  respectively lie on the circle with diameter  $OL$ , and the line  $MN$  passes through  $I$ . Hence we can construct the point  $N$  and the diagonal  $BD$ .

8. (8–9, K.Belskij) Let  $ABCD$  be a quadrilateral with  $\angle B = \angle D$  and  $AD = CD$ . The incircle of triangle  $ABC$  touches the sides  $BC$  and  $AB$  at points  $E$  and  $F$  respectively. Prove that the midpoints of segments  $AC, BD, AE$ , and  $CF$  are concyclic.

**Solution.** Let  $K, L, M, N$  be the midpoints of  $BD, AC, AE, CF$  respectively. Since  $LM \parallel BC$ , and  $LN \parallel AB$ , we have  $\angle MLN = \angle CBA$ . On the other hand  $\overrightarrow{KM} = (\overrightarrow{DA} + \overrightarrow{BE})/2$ ,  $\overrightarrow{KN} = (\overrightarrow{DC} + \overrightarrow{BF})/2$ , and since  $DA = DC, BE = BF$ , and the angle between  $\overrightarrow{DA}$  and  $\overrightarrow{DC}$  equals the angle between  $\overrightarrow{BE}$  and  $\overrightarrow{BF}$ , we obtain that the angle  $MKN$  also is equal to these angles.

9. (8–9, A.Mardanov) Let  $ABCD$  ( $AD \parallel BC$ ) be a trapezoid circumscribed around a circle  $\omega$ , which touches the sides  $AB, BC, CD$ , and  $AD$  at points  $P, Q, R, S$  respectively. The line passing through  $P$  and parallel to the bases of the trapezoid meets  $QR$  at point  $X$ . Prove that  $AB, QS$ , and  $DX$  concur.

**First solution.** Let  $I$  be the center of  $\omega$ . Then the sidelines of triangles  $PQX$  and  $AID$  are parallel (fig. 9). Thus these triangles are homothetic which yields the required concurrency.




Fig. 9

**Second solution.** Fix the points  $A, B, P, Q, S$  and move  $R$  along  $\omega$ . Then  $D$  and  $X$  move along  $AS$  and parallel line passing through  $P$  respectively. Clearly the correspondence between  $D$  and  $X$  is projective, and since they meet at infinity, this correspondence is linear, i.e. all lines  $DX$  are concurrent. Also it is clear that their common point lies on  $AB$ , hence it is sufficient to find one position such that  $DX, AB$ , and  $QS$  concur. This is correct if the trapezoid  $ABCD$  is isosceles.

10. (8–9, A.Tereshin) Let  $\omega$  be the circumcircle of a triangle  $ABC$ . A point  $T$  on the line  $BC$  is such that  $AT$  touches  $\omega$ . The bisector of angle  $BAC$  meets  $BC$  and  $\omega$  at points  $L$  and  $A_0$  respectively. The line  $TA_0$  meets  $\omega$  at point  $P$ . The point  $K$  lies on the segment  $BC$  in such a way that  $BL = CK$ . Prove that  $\angle BAP = \angle CAK$ .

**Solution.** The projection of  $\omega$  to itself from  $T$  swaps  $B$  with  $C$ ,  $P$  with  $A_0$ , and conserve  $A$ . This yields the equality of cross-ratios  $(BCA_0P) = (CBAP)$ , i.e.  $\sin \angle BAP : \sin \angle CAP = PB : PC = AB^2 : AC^2$ . On the other hand applying the sines law to triangles  $AKC$  and  $BKC$  we obtain that  $\sin \angle CAK : \sin \angle BAK = (CK/AC) : (BK/AB) = (AB/AC) \cdot (BL/AL) = AB^2 : AC^2$ , which yields the required equality.

11. (8–10, B.Butyrin) Let  $M, N$  be the midpoints of sides  $AB, AC$  respectively of a triangle  $ABC$ . The perpendicular bisector to the bisectrix  $AL$  meets the bisectrices of angles  $B$  and  $C$  at points  $P$  and  $Q$  respectively. Prove that the common point of lines  $PM$  and  $QN$  lies on the tangent to the circumcircle of  $ABC$  at  $A$ .

**Solution.** Note that the lines  $PQ$  and  $MN$  meet at the midpoint  $K$  of segment  $AL$ . Also  $P$  bisects the arc  $AL$  of circle  $ABL$ , therefore  $\angle BPL = \angle CAL = \angle BIC - \pi/2$ , where  $I$  is the incenter of  $ABC$ , i.e.  $PL \perp CI$ . Similarly  $QL \perp BI$ . Thus the sidelines of triangle  $PQL$  are parallel to the sidelines of triangle formed by the touching points of sides of  $ABC$  with the incircle, and the tangents to the circumcircle of  $PQL$  at the vertices are parallel to the sidelines of  $ABC$ . Since the circle  $APQ$  is the reflection of circle  $LPQ$  about  $PQ$ , the tangent to this circle at  $P$  is parallel to  $AB$ , and the tangent at  $A$  coincide with the tangent to the circumcircle of  $ABC$ . Also  $\angle PAQ = \angle PLQ = \pi - \angle PIQ$ , hence  $I$  lies on the circle  $APQ$ , and the tangent to this circle at  $I$  is parallel to  $BC$ . Denote the common point of tangents at  $A$  and  $I$  as  $T$  (fig. 11)




Fig. 11

We have to prove that  $PM$  passes through  $T$ , i.e. that the line passing through  $A$  and parallel to the tangent at  $P$ , and the line passing through the projection  $K$  of  $P$  to  $AI$ , parallel to the tangent at  $I$  meet  $AT$  at the same point. Let  $S$  be the common point of tangents at  $A$  and  $P$ ,  $U$  be the common point of  $AI$  with the line passing through  $P$  and parallel to  $IT$ . then  $IK : KU = AT : AS = \operatorname{ctg} \angle IPA : \operatorname{ctg} \angle AIP$ , and we obtain the required assertion.

12. (8–10, D.Shvetsov) The bisectors  $AA_1, CC_1$  of a triangle  $ABC$  with  $\angle B = 60^\circ$  meet at point  $I$ . The circumcircles of triangles  $ABC, A_1IC_1$  meet at point  $P$ . Prove that the line  $PI$  bisects the side  $AC$ .

**Solution.** Since  $\angle A_1IC_1 = 120^\circ = 180^\circ - \angle A_1BC_1$ , the circles  $ABC$  and  $A_1IC_1$  meet at points  $B$  and  $P$  (fig. 12). Hence the triangles  $PA_1C$  and  $PC_1A$  are similar, i.e.  $PA_1 : PC_1 = A_1C : AC_1$ . On the other hand since  $\angle AC_1I + \angle IA_1C = 180^\circ$ , we obtain applying the sines law to the triangles  $AC_1I$  and  $CA_1I$  that  $A_1C : AC_1 = IC : IA$ . therefore  $\sin \angle PIA_1 : \sin \angle PIC_1 = IC : IA$ , and  $IP$  is the median of triangle  $IAC$ .




Fig. 12.

13. (8–11, A.Zaslavsky) Can an arbitrary polygon be cut into isosceles trapezoids?

**Answer.** Yes.

**Solution.** Since an arbitrary polygon can be cut into triangles, and an arbitrary triangle can be cut into isosceles triangles (drawing the altitude to the longer side and joining its foot with the midpoints of two remaining sides), It is sufficient to solve the problem for isosceles triangles.

Note that we can cut an isosceles triangle into three isosceles trapezoids drawing three rays parallel to its sides from the center (fig. 13.1).




Fig. 13.1

Now if the angle at the vertex of the triangle is greater than  $60^\circ$  we can cut it into isosceles trapezoids applying several times the construction on fig. 13.2.




Fig. 13.2

Finally if the angle at the vertex of the triangle is less than  $60^\circ$  cut it into three triangles joining the vertices with the circumcenter. Two of obtained triangles are obtuse-angled, and the angle at the vertex of third one is twice greater than the angle of the original triangle. Repeating this trick several times we cut the given triangle into isosceles triangles with the angles at the vertices greater than  $60^\circ$ .

14. (9–11, A.Tereshin) The incircle  $\omega$  of a right-angled triangle  $ABC$  touches the circumcircle of its medial triangle at point  $F$ . Let  $OE$  be the tangent to  $\omega$  from the midpoint  $O$  of the hypotenuse  $AB$ , distinct from  $AB$ . Prove that  $CE = CF$ .



**Solution.** The homothety with center  $C$  and coefficient 2 maps  $F$  to the touching point of the circumcircle and the semiincircle of given triangle. Hence the line  $CF$  is the reflection about the bisector of angle  $C$  of the line joining  $C$  with the touching point of the hypotenuse with the corresponding excircle. Let the hypotenuse touches the incircle and the excircle at points  $T$  and  $S$  respectively. Since  $OE = OT = OS$  we obtain that  $\angle SET = \pi/2$ , i.e. the line  $SE$  passes through the point of  $\omega$  opposite to  $T$ . But  $SC$  also passes through this point, therefore  $E$  lies on  $SC$  and is the reflection of  $F$  about the bisector of angle  $C$  (fig. 14).




Fig. 14

15. (9–11, M.Panov) The difference of two angles of a triangle is greater than  $90^\circ$ . Prove that the ratio of its circumradius and inradius is greater than 4.

**First solution.** Let  $A$  be the smallest angle of triangle  $ABC$ ,  $B$  be the greatest angle,  $O$  be the circumcenter,  $L$  be the midpoint of arc  $AB$ ,  $CD$  and  $PQ$  be the chord and the diameter of the circumcircle parallel to  $AB$  (fig. 15). Since  $\sphericalangle CD = \sphericalangle ADC - \sphericalangle AD = \sphericalangle ADC - \sphericalangle BC > \pi$ , we obtain that  $A, B, C$  lie on the same semiplane with respect to  $PQ$ , i.e.  $\angle OLC > \pi/4$ , and the distance from  $O$  to the line  $LC$  passing through the incenter  $I$  is greater than  $R/\sqrt{2}$ . Therefore  $OI^2 = R^2 - 4Rr > R^2/2$  which is equivalent to the required inequality.




Fig. 15

**Second solution.** From the formula  $r = 4R \sin(A/2) \sin(B/2) \sin(C/2)$  we have

$$\frac{r}{R} = 2 \sin \frac{C}{2} \left( \cos \frac{B-A}{2} - \cos \frac{A+B}{2} \right) < 2 \sin \frac{C}{2} \left( \frac{1}{\sqrt{2}} - \sin \frac{C}{2} \right) \leq 2 \left( \frac{1}{2\sqrt{2}} \right)^2 = \frac{1}{4}.$$

**Remark.** the obtained estimation is exact. If  $\cos C = 3/4$ ,  $A = \pi/4 - C/2$ ,  $B = 3\pi/4 - C/2$  all inequalities transform to equalities and  $R = 4r$ .

16. (9–11, A.Mardanov) Let  $AA_1$ ,  $BB_1$ , and  $CC_1$  be the bisectors of a triangle  $ABC$ . The segments  $BB_1$  and  $A_1C_1$  meet at point  $D$ . Let  $E$  be the projection of  $D$  to  $AC$ . Points  $P$  and  $Q$  on the sides  $AB$  and  $BC$  respectively are such that  $EP = PD$ ,  $EQ = QD$ . Prove that  $\angle PDB_1 = \angle EDQ$ .

**Solution.** The sum of distances from any point of segment  $A_1C_1$  to  $AB$  and  $BC$  equals to the distance from this point to  $AC$  (because this is correct for the endpoints of the segment). Since the distances from  $D$  to  $AB$  and  $BC$  are equal each of these distances equals to a half of  $DE$ , i.e.  $D$  is the incenter of triangle  $BPQ$  (fig. 16). Thus  $\angle EPQ = \angle DPB$ , and  $\angle EQP = \angle DQB$ . Therefore  $B$  and  $E$  are isogonally conjugated with respect to triangle  $DPQ$  and we obtain the required equality.




Fig. 16

17. (9–11, L.Dong) Let  $ABC$  be a non-isosceles triangle,  $\omega$  be its incircle. Let  $D, E$  and  $F$  be the points at which the incircle of  $ABC$  touches the sides  $BC, CA$  and  $AB$ , respectively. Let  $M$  be the point on ray  $EF$  such that  $EM = AB$ . Let  $N$  be the point on ray  $FE$  such that  $FN = AC$ . Let the circumcircles of  $\triangle BFM$  and  $\triangle CEN$  intersect  $\omega$  again at  $S$  and  $T$ , respectively. Prove that  $BS, CT$  and  $AD$  concur.

**Solution.**

**Lemma.** Let  $X, Y, Z$  be points on  $\omega$  such that  $DX, EY, FZ$  concur. Then  $AX, BY, CZ$  concur.

**Proof.** Clearly,  $\angle AEX = \angle EDX$  и  $\angle XFA = \angle XDF$ . Also by the sines law

$$\frac{\sin \angle FAX}{\sin \angle XFA} = \frac{XF}{AX}, \quad \frac{\sin \angle XAE}{\sin \angle AEX} = \frac{XE}{AX}.$$

Therefore

$$\frac{\sin \angle BAX}{\sin \angle XAC} = \frac{\sin \angle FAX}{\sin \angle XAE} = \frac{XF}{XE} \cdot \frac{\sin \angle XFA}{\sin \angle AEX} = \frac{XF}{XE} \cdot \frac{\sin \angle XDF}{\sin \angle EDX} = \left( \frac{\sin \angle XDF}{\sin \angle EDX} \right)^2.$$

Similarly

$$\frac{\sin \angle CBY}{\sin \angle YBA} = \left( \frac{\sin \angle YED}{\sin \angle FEY} \right)^2; \quad \frac{\sin \angle ACZ}{\sin \angle ZCB} = \left( \frac{\sin \angle ZFE}{\sin \angle DFZ} \right)^2.$$

Hence

$$\frac{\sin \angle BAX}{\sin \angle XAC} \cdot \frac{\sin \angle CBY}{\sin \angle YBA} \cdot \frac{\sin \angle ACZ}{\sin \angle ZCB} = 1.$$

and the assertion of the lemma follows from the Ceva theorem.

Return to the problem.

Denote by  $O(ABCD)$  the cross ratio of lines  $OA, OB, OC, OD$ . Let  $J$  be the common point of  $FT$  and  $ES$ ,  $G$  be the second common point of  $AD$  and  $\omega$ ,  $K$  be the common point of  $EF$  and  $BC$  (fig. 17).




Fig. 17

Since  $\angle FMS = \angle FBS$ ,  $\angle MES = \angle BFS$ , we obtain that the triangles  $SBF$  and  $SME$  are similar. Thus  $SE : SF = ME : BF = AB : BF$ . Then

$$E(AFJD) = E(EFSD) = (EFSD) = \frac{SE}{SF} : \frac{DE}{DF} = \frac{AB}{BF} \cdot \frac{DF}{DE}.$$

Similarly

$$F(AEJD) = \frac{AC}{CE} \cdot \frac{DE}{DF}.$$

And so

$$E(AFJD) : F(AEJD) = \frac{AB}{AC} \cdot \frac{CE}{BF} \cdot \frac{DF^2}{DE^2} = \frac{AB}{AC} \cdot \frac{CE}{BF} \cdot \frac{KF}{KE}.$$

Applying the Menelaos theorem to the triangle  $AEF$  and points  $K, B, C$ , we obtain

$$\frac{AB}{BF} \cdot \frac{FK}{KE} \cdot \frac{EC}{CA} = 1.$$

Therefore  $E(AFJD) = F(AEJD)$ , i.e.  $A, J, D$  are collinear. Hence  $DG, ES$ , and  $FT$  concur at  $J$ , and by the lemma  $BS, CT$ , and  $AG$  concur.

18. (9–11, D.Shvetsov) Let  $AA_1, BB_1, CC_1$  be the altitudes of an acute-angled triangle  $ABC$ ;  $I_a$  be its excenter corresponding to  $A$ ;  $I'_a$  be the reflection of  $I_a$  about the line  $AA_1$ . Points  $I'_b, I'_c$  are defined similarly. Prove that the lines  $A_1I'_a, B_1I'_b, C_1I'_c$  concur.

**First solution.** The lines  $A_1A, A_1B, A_1A'$ , and  $A_1I_a$  form a harmonic quadruple. Therefore their meeting points with  $AL$  also form a harmonic quadruple. Since the fourth harmonic point for  $A, L, I_a$  coincide with the incenter  $I$  of triangle  $ABC$ , we obtain that  $A_1A'$  passes through  $I$ . Similarly  $B_1B'$  and  $C_1C'$  pass through  $I$ .

this reasoning can be modified. Since  $A_1A, A_1B, A_1I$ , and  $A_1I_a$  form a harmonic quadruple, and  $A_1A \perp A_1B$ , the lines  $A_1A$  and  $A_1B$  bisect the angles between  $A_1I$  and  $A_1I_a$ . Therefore  $A_1A'$  passes through  $I$ .

**Second solution.** Prove that  $A_1I_a, B_1I_b, C_1I_c$  concur. then their reflections  $A_1A', B_1B', C_1C'$  about the bisectors of triangle  $A_1B_1C_1$  also concur. Note that for example  $\sin \angle I_b I_a A_1 : \sin \angle I_c I_a A_1 = (BA_1 : CA_1) \cdot (I_a C : I_a B)$ . Applying the Ceva theorem to the lines  $I_a A, I_b B, I_c C$  and  $AA_1, BB_1, CC_1$  we obtain that the product of this ratio and two similar ones equals 1.

**Remark.** The assertion of the problem is a partial case of the following fact. If points  $A_1, B_1, C_1$  lie on the sidelines  $BC, CA, AB$  of triangle  $ABC$ , and points  $A_2, B_2, C_2$  lie on the sidelines  $B_1C_1, C_1A_1, A_1B_1$  of triangle  $A_1B_1C_1$ , in such a way that  $AA_1, BB_1, CC_1$  concur, and  $A_1A_2, B_1B_2, C_1C_2$  concur, then  $AA_2, BB_2, CC_2$  also concur.

19. (10–11, M.Evdokimov) A triangle  $ABC$ , its circumcircle, and its incenter  $I$  are drawn on the plane. Construct the circumcenter of  $ABC$  using only a ruler.

**Solution.** Construct the common point  $C_1$  of tangents to the circle at points  $A, B$  and the second common point  $C_2$  of the circle and the line  $CI$ . The line  $C_1C_2$  is the perpendicular bisector to the segment  $AB$  therefore it passes through the circumcenter. Constructing similarly the perpendicular bisector to  $AC$  find the circumcenter.

20. (10–11, L.Shatunov) Lines  $a_1, b_1, c_1$  pass through the vertices  $A, B, C$  respectively of a triangle  $ABC$ ;  $a_2, b_2, c_2$  are the reflections of  $a_1, b_1, c_1$  about the corresponding bisectors of  $ABC$ ;  $A_1 = b_1 \cap c_1, B_1 = a_1 \cap c_1, C_1 = a_1 \cap b_1$ , and  $A_2, B_2, C_2$  are defined similarly. Prove that the triangles  $A_1B_1C_1$  and  $A_2B_2C_2$  have the same ratios of the area and the circumradius (i.e.  $\frac{S_1}{R_1} = \frac{S_2}{R_2}$ , where  $S_i = S(\Delta A_i B_i C_i), R_i = R(\Delta A_i B_i C_i)$ ).

**Solution.**

**Lemma.** Let  $X', Y', Z'$  lie on the sides  $YZ, ZX, XY$  respectively of triangle  $XYZ$ . Then

$$S_{X'Y'Z'} = \frac{XY' \cdot YZ' \cdot ZX' + X'Y \cdot Y'Z \cdot Z'X}{4R_{XYZ}}.$$

(the points  $X', Y', Z'$  may also lie on the extensions of the sides. In this case we have to consider the segments in the formula as oriented.)

**Proof.** Let  $XY' = \alpha XZ, YZ' = \beta YX, ZX' = \gamma ZY$ . Then  $S_{X'Y'Z'} : S_{XYZ} = 1 - \alpha(1 - \beta) - \beta(1 - \gamma) - \gamma(1 - \alpha) = \alpha\beta\gamma + (1 - \alpha)(1 - \beta)(1 - \gamma)$  and the assertion of the lemma follows from the formula  $S_{XYZ} = (XY \cdot YZ \cdot ZX)/4R_{XYZ}$ .

Now apply the lemma to the triangle  $A_1B_1C_1$  and  $A, B, C$  on its sidelines. Denoting  $\angle B_1AC = \alpha, \angle C_1BA = \beta, \angle A_1CB = \gamma$  we obtain (using the sines law for the triangles  $AB_1C_1, BC_1A_1, CA_1B_1$ )

$$S_{ABC} = \frac{AB \cdot BC \cdot CA(\sin \alpha \sin \beta \sin \gamma + \sin(A + \alpha) \sin(B + \beta) \sin(C + \gamma))}{4R_1 \sin \angle A_1 B_1 C_1 \sin \angle B_1 C_1 A_1 \sin \angle C_1 A_1 B_1}.$$

Applying the sines law to the triangle  $A_1B_1C_1$  we obtain that the denominator equals  $2S_1/R_1$ . Finally note that if we replace the triangle  $A_1B_1C_1$  to  $A_2B_2C_2$  the numerator do not change.

21. (10–11, A.Zaslavsky) A chord  $PQ$  of the circumcircle of a triangle  $ABC$  meets the sides  $BC, AC$  at points  $A', B'$  respectively. The tangents to the circumcircle at  $A$  and  $B$  meet at point  $X$ , and the tangents at points  $P$  and  $Q$  meet at point  $Y$ . The line  $XY$  meets  $AB$  at point  $C'$ . Prove that the lines  $AA', BB',$  and  $CC'$  concur.

**Solution.** Let  $PQ$  and  $AB$  meet at point  $U, AA'$  and  $BB'$  meet at point  $V$ . Then the line  $XY$  is the polar of  $U$  with respect to the circumcircle, therefore  $A, B, U, C'$  form a harmonic quadruple. Thus  $CV$  passes through  $C'$ .

22. (10–11, D.Reznik, A.Akopyan) A segment  $AB$  is given. Let  $C$  be an arbitrary point of the perpendicular bisector to  $AB$ ;  $O$  be the point on the circumcircle of  $ABC$  opposite to  $C$ ; and an ellipse centered at  $O$  touch  $AB, BC, CA$ . Find the locus of touching points of the ellipse with the line  $BC$ .

**Answer.** The circle with diameter  $BQ$ , where  $Q$  is a point on the ray  $AB$  such that  $AQ = 3AB/2$ , without  $B$  and  $Q$ .

**Solution.** Let  $M$  be the midpoint of  $AB, N$  be the reflection of  $M$  about  $O, P$  be the common point of  $PN$  and  $BC, U, V$  be the common points of the line passing through  $N$  and parallel to  $AB$  with  $BC, AC$  respectively (fig. 22). Since the triangles  $BPQ$  and  $UPN$  are similar we have  $BP : PN = BQ : UN = AB : UV$ , i.e. the line passing through  $P$  and parallel to  $AB$  passes through the common point of diagonals of trapezoid  $ABUV$ . Therefore the ellipse inscribed into the trapezoid touches  $BC$  at point  $P$ . Since

$PN \parallel OB \perp BC$  this point lies on the circle with diameter  $BQ$ . It is clear that all points of this circle distinct from  $B$  and  $Q$  belong to the required locus.




Fig. 22

23. (10–11, I.Kukharchuk) A point  $P$  moves along a circle  $\Omega$ . Let  $A$  and  $B$  be fixed points of  $\Omega$ , and  $C$  be an arbitrary point inside  $\Omega$ . The common external tangents to the circumcircles of triangles  $APC$  and  $BCP$  meet at point  $Q$ . Prove that all points  $Q$  lie on two fixed lines.

**Solution.** The common point of external tangents is the center of circle  $\omega$  such that the inversion with respect to it swaps the circles  $APC$  and  $BPC$ . Consider an inversion centered at  $C$  transforming  $A, B, P$  to  $A', B', C'$  respectively. It maps the circles  $APC, BPC$  to the lines  $A'P', B'P'$  respectively, the image of  $\omega$  is the bisector of some angle between these lines, and the image of  $Q$  is the reflection  $Q'$  of  $C$  about this bisector. Since the bisectors of angles between  $P'A'$  and  $P'B'$  pass through two fixed points — the midpoints of arcs  $A'B'$  of circle  $A'B'P'$ , all points  $Q'$  lie on two circles centered at these points and passing through  $C$ . The considered inversion maps these circles to two fixed lines.

**Remark.** The point  $Q$  jumps from one line to the second one when  $P$  intersects one of lines  $AC, BC$ .

24. (11, Tran Quang Hung, N.Dergiados) Let  $SABC$  be a pyramid with right angles at the vertex  $S$ . Points  $A', B', C'$  lie on the edges  $SA, SB, SC$  respectively in such a way that the triangles  $ABC$  and  $A'B'C'$  are similar. Does this yield that the planes  $ABC$  and  $A'B'C'$  are parallel?

**Answer.** Yes.

**First solution.** Suppose that the planes  $ABC$  and  $A'B'C'$  are not parallel. Then apply the homothety centered at  $S$  mapping  $A'B'C'$  to a triangle congruent to  $ABC$ , and

transform the obtained triangle to  $ABC$  by an isometry of the space. The image of  $S$  is a point  $S'$  distinct from  $S$  and its reflection about the plane  $ABC$ . On the other hand both points  $S, S'$  lie on three spheres with diameters  $AB, BC, CA$ . Since the centers of these spheres are not collinear they have only two common points symmetric with respect to the plane  $ABC$  — contradiction.

**Second solution.** Let  $A'B' = tAB$ . Then  $B'C' = tBC, C'A' = tAC$ . Since the angles at  $S$  are right we have  $SA'^2 + SB'^2 = t^2AB^2, SA'^2 + SC'^2 = t^2AC^2, SB'^2 + SC'^2 = t^2BC^2$ . From this we obtain that  $SA'^2 = t^2(AB^2 + AC^2 - BC^2)/2 = t^2SA^2$ , i.e.  $SA' = tSA$ . Similarly  $SB' = tSB, SC' = tSC$  and therefore the planes  $ABC$  and  $A'B'C'$  are parallel.